



Fundacja na rzecz Nauki Polskiej

Quasars as tracers of cosmic flows

Bożena Czerny

Copernicus Astronomical Center, Warsaw

In collaboration with

M. Bilicki, M. Chodorowski, K. Hryniewicz, M. Krupa, J. Modzelewska,
F. Petrogalli, A. Pollo, W. Pych, A. Swieton, A. Udalski, Weimin Yuan

Versoix, 18 March 2014


Role of quasars in cosmology

- Active role: quasars affect the galaxy formation
- Passive role: quasars can be used to trace the expansion history of the Universe

Quasars – their place in the Universe


- Quasars are distant/bright subclass of Active Galactic Nuclei
- Only 1 in a million of normal galaxies is a quasar but all galaxies went through some AGN activity stage at some point
- AGN activity is a non-stellar radiation due to material generously falling into the central black hole

AGN schematic structure




~ Urry & Padovani 1995

Impressive career of AGN



“They are no ornament which just happens to be there...” A. Fabian

Turning point I: 1998




Magorrian et al. :

$$M_{\text{bh}} = 0.0052 M_{\text{bulge}}$$

It implies the existence of the mechanism regulating the common growth of a galaxy and a central black hole.

Nowadays mostly in M-sigma relation form.

Current version of the plot:




Schramm & Silverman 2013

Central black hole and host galaxy coexistence

- Massive central black holes are in all galaxies
- In most galaxies the activity is low (e.g. Sgr A* in the Milky Way) but there is no strict border between AGN and non-active galaxies
- Thus again BH and galaxies likely evolve together

Turning point II: 1998




Boyle and Trelevich (1998) - the plot of the cosmic evolution of quasars and Star Formation Rate (SFR).

Also Richstone et al.98




Implies that quasar activity and star formation proceed together!

Most recent version of quasar-SFR coevolution plot




Zheng et al. 2009

Large scale structure formation needs quasars



S. White talk

Cosmological simulations need AGN input



Simulations by McCarthy et al. based on OverWhelmingly Large Simulations project. Without (REF) and with AGN input.

Black histogram: data from Loubser et al. 2009

Other aspects:

- galaxies increase in size with time but their masses do not grow – AGN induced puffs up (Ishibashi et al. 2013)
- Too many stars
- Wrong shape of luminosity function
- CD galaxies too massive...

In general, the AGN feedback is required:



- In clusters of galaxies (to stop the efficient cooling flows)
- In groups of galaxies (to control the group structure through regulation of gas mass fraction)
- In individual galaxies to control the star formation and cold gas and provide M-sigma relation

Quasars as a tool in cosmology

- Quasars are numerous
- Quasars are seen up to redshift 7
- Quasars do not show significant evolution with redshift in their properties

**SO WHY NOT TO USE THEM TO TRACE
THE EXPANSION OF THE UNIVERSE AND
THE DARK ENERGY ?**

The largest puzzle of cosmology: Universe content




Recent results from Planck:

These 4.9 % of barionic matter consists of:

Stars - 0.5 %
Gas - 4.1 %
Neutrinos - 0.3 %

We need more independent tests to believe!

Measuring the expansion of the Universe with a single object



Hubble diagram

Measuring the expansion of the Universe with a single object





FIGURE 1


Velocity-Distance Relation among Extra-Galactic Nebulae.

Original Hubble paper

Measuring the expansion of the Universe with a single object



Hubble diagram




Usual diagram for SN Ia

Luminosity distance

$$D_L = \frac{L_{\text{intr}}}{4 \pi F_{\text{obs}}} \quad 1/2$$

So the problem reduces to determination of the absolute luminosity NOT from the redshift but independently

In SN Ia this happens since SN Ia are 'standard candles'



<https://www.llnl.gov/str/SepOct08/hoffman.html>


Now we have to do 'the same' for quasars

Absolute luminosity.I.

Dusty
Molecular
Torus

Hryniewicz 2013


Lines are
delayed
with
respect to
continuum !



Broad
Line
Region

Accretion
Disc

Black
Hole



Absolute luminosity.II.




Fig. 1. The BLR region covers the range of the disk with an effective temperature lower than 1000 K: the dusty wind rises and then breaks down when exposed to the radiation from the central source. The dusty torus is the disk range where the irradiation does not destroy the dust and the wind flows out.

Theory outlined in Czerny & Hryniewicz (2011):

- Large outflow forms in the region where the disk temperature is below 1000 K and allows for dust formation
- Outflow is caused by radiation pressure acting on dust grains
- Far from the disk the dusty clouds are irradiated and dust evaporates
- Dustless material loses support against gravity and falls back
- Failed wind forms

Absolute luminosity.III.

From Shakura-Sunyaev accretion disk model the disk temperature, T_{eff} , as a function of disk radius is

$$F(R) = \sigma T_{eff}^4 = \frac{3GM\dot{M}}{8\pi R^3} \quad (1)$$

i.e. assuming that BLR starts at $T_{eff} = 1000K$, where dust forms in the disk atmosphere we can obtain this radius $R = R_{BLR}$

$$R_{BLR} = \frac{3GM\dot{M}}{8\pi\sigma(1000K)^4}^{1/3} \quad (2)$$

i.e. depends on the black hole mass, M and accretion rate, \dot{M} . But again from Shakura-Sunyaev accretion disk model the monochromatic luminosity from the whole disk at a fixed frequency is given by

$$L_\nu = 0.91 \frac{\nu}{10^{15}Hz}^{1/3} (M\dot{M})^{2/3} \cos i \quad (3)$$

This gives the know trend with frequency, $L_\nu \propto \nu^{1/3}$, and for a fixed frequency, ν , combined with Eq.* reproduces the result known from reverberation of nearby active galaxies

$$R_{BLR} = const \quad L_\nu^{1/2} \quad (4)$$

const contains only known physical/mathematical constants, does not depend on M or \dot{M} .

Lines good for reverberation in the optical band

Proper lines, to be consistent with our BLR picture, should not show net outflow

Hbeta - nearby objects

Mg II - intermediate quasars

CIV - high redshift quasars ?

This line does not belong to LIL

Spectroscopic studies of time delay of line vs. continuum

THE ASTROPHYSICAL JOURNAL LETTERS, 740:L49 (5pp), 2011 October 20

WATSON ET AL.




Figure 2. AGN Hubble diagram. The luminosity distance indicator τ/\sqrt{F} is plotted as a function of redshift for 38 AGNs with H β lag measurements. On the right axis the luminosity distance and distance modulus ($m - M$) are shown using the SBF distance to NGC 3227 as a calibrator. The current best cosmology (Komatsu et al. 2011) is plotted as a solid line. The line is not fit to the data but clearly follows the data well. Cosmologies with no dark energy components are plotted as dashed and dotted lines. The lower panel shows the logarithm of the ratio of the data compared to the current cosmology on the left axis, with the same values but in magnitudes on the right. The red arrow indicates the correction for internal extinction for NGC 3516. The green arrow shows where NGC 7469 would lie using the revised lag estimate from Zu et al. (2011). NGC 7469 is our largest outlier and is believed to be an example of an object with a misidentified lag (Peterson 2010).

(A color version of this figure is available in the online journal.)

Past reverberation studies:

- about 40 nearby AGN
- about 10 $z < 0.4$ quasars
- 7 distant objects but only 1 detection

$$R_{BLR} = \text{const} \quad L_{\nu}^{1/2}$$

Watson et al. 2011

Spectroscopic studies of time delay of line vs. continuum

$$R_{BLR} = \text{const} \quad L_{\nu}^{1/2}$$

Our SALT current campaign:
Three $z = 1$ quasars

SALT – Southern African Large Telescope

Poland has 10 % SAL time.



SALT




Construction similar
to Hobby-Eberly
Telescope

Limited movement,
difficult calibration.

Size: 10-11 m

91 mirrors, 1.2 m
each


Exemplary SALT spectra



Spectra are nice but we need about 5 spectra per year, for 5 years, to measure the expected delay of about 500 days

Hryniewicz et al. 2014

Exemplary SALT spectra



That one does not make us happy

Spectroscopic studies: alternative to SALT





Advantages and disadvantages:

- A few thousands of quasars in comparison with three in SALT
- Much lower accuracy; 6 – 10 percent error in line measurement in comparison with 1 percent in SALT

LAMOST spectroscopic survey telescope in China

Pure photometric multi-channel studies

We also simulate now the possibilities to use the future LSST (Large Synoptic Sky Survey). Preliminary estimates of the line contribution to the photometric channels.



Other ideas to use quasars to trace cosmic expansion

- Selection of the exactly Eddington rate quasars (Marziani & Sulentic 2013)
- Doing reverberation by the dusty torus in the optical and IR (Hoenig 2014; Yoshii et al. 2014)

First quasar with tentatively measured delay from HET




Fig. 8. *Top:* Hubble diagram of the combined sample. The distance modulus redshift relation of the best-fit Λ CDM cosmology for a fixed $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ is shown as the black line. *Bottom:* Residuals from the best-fit Λ CDM cosmology as a function of redshift. The weighted average of the residuals in logarithmic redshift bins of width $\Delta z/z \sim 0.24$ are shown as black dots.

Quasar S5 0836+7

HET 7 yr monitoring
Kaspi et al. (2007)

$\tau_{\text{rest}} = 595 \text{ days}$

$Z = 2.172$

$\Delta \mu = + 0.39$

Figure from Betoule et al.
2014

Summary

- Quasars are important ingredients of the Universe
- Quasars can be used to determine the expansion rate of the Universe, i.e. the cosmological model and the dark energy properties
- They will describe – but not explain – the dark energy nature but better quantitative measurements will provide good starting point
- So far, cosmological constant provides good enough description...