

The impact of 1-m class telescopes on our understanding of classical pulsators' dynamics

R. Smolec

Nicolaus Copernicus Astronomical Center, Warsaw, Poland; e-mail: smolec@camk.edu.pl

Observing projects with 1-m class telescopes

Optical Gravitational Lensing Experiment, OGLE, is a long-term photometric sky survey led by Warsaw University astronomers (Udalski et al., 1992). OGLE observes with 1.2-m telescope located in Las Campanas Observatory in Chile (Fig. 1, left; Fig. 2). It regularly monitors hundreds of fields in the direction of Galactic bulge, Galactic disc and the Magellanic System. Although its original goal was the detection of gravitational lensing events, with its top-quality and regular observations OGLE contributed to nearly all fields of astrophysics. In particular, it gathered top-quality photometry for thousands of classical pulsators: Cepheids and RR Lyrae stars that revolutionized our knowledge about these variables. Read more at http://ogle.astrouw.edu.pl/.

Cluster AgeS Experiment, CASE (e.g. Kaluzny et al. 2005) is the long-term projects that observes the globular clusters with the goal of determining the clusters' characteristics (eg., age, metallicity) by searching and analysing the variable stars. It frequently uses the 1-m Swope telescope (Fig. 1, right) located at Las Campanas observatory that was also used in the first phase of the OGLE project. CASE collected top-quality, long-baseline photometry for tens of RR Lyr stars in a few globular clusters. Read more at http://case.camk.edu.pl

Petersen Diagram Revolution



A few years ago, only double-mode fundamental mode (F) and first overtone (10) RR Lyr stars were known (RRd stars, black dots in Fig. 5). Then, a few F+2O stars were discovered, mostly among the Blazhko (modulated) variables (filled green triangles). For the discovery or for understanding of the other classes marked in Fig. 5, exploration of the OGLE data was crucial. These new classes of double-periodic variables are the following:

* $RR_{0.61}$ stars, with radial 1O and non-radial $\ell = 8$ and 9 modes excited (small red squares). Most of the known stars were discov-ered in the OGLE data (Netzel, Smolec & Moskalik, 2015a,b). \star $\mathrm{RR}_{0.68}$ stars (magenta diamonds), with radial 1O and additional,

long-period variability excited. Discovered in the OGLE data (Netzel, Smolec & Dziembowski, 2015).

* Anomalous RRd stars (open blue circles). F+1O pulsators with anomalous period and amplitude ratios and frequent modulation of the excited modes. Discovered in the OGLE data (e.g., Soszyński et al., 2016)

* Extreme RRd stars (red open squares). F+1O pulsators with extremely long periods. Discovered in the OGLE data (Smolec et al., 2016)

 \star Double-periodic stars with a typical light curve shape, in between the shapes characteristic for F-mode and 10 mode pulsators (open green triangles). Discovered in the OGLE data (Prudil et al., 2017).

Kulterickes Kaluzny J., et al. 2005, AIP Conf. Proc., **752**, 70 Netzel H., Smolec R., Moskalik P., 2015a, MNRAS, **447**, 1173 Netzel H., Smolec R., Moskalik P., 2015b, MNRAS, **453**, 2022 Netzel H., Smolec R., Dzienbowski W. 2015, MNRAS, **451**, L25

References

Prudil Z. et al., 2017, MNRAS, **4074**, 2349 Smolec R., 2017, MNRAS, **468**, 4299 Smolec R. et al., 2016, MNRAS, **461**, 2934 Smolec R. et al., 2016, MNRAS, **467**, 2349 Soszyński I. et al., 2016, MNRAS, **467**, 233 Udalski A. et al., 1992, AcA, **42**, 253



The 1.2-m OGLE (left) and 1.0-m Swop



Fig. 2. The OGLE telescope in Las Ca ares 1 & 2 from http://ogle.astrouw.edu.pl/ and





In 51 fundamental mode Cepheids of the OGLE collection pe riodic modulation of the light curves was detected (Smolec, 2017; Fig. 6). Although the overall incidence rate is very low, $\approx 1\%$ in each of the Magellanic Clouds, in the case of the SMC and pulsation periods between 12 and 16 d the incidence rate is nearly 40%. Typical modulation periods are between 70 and 300 d. In nearly all stars the mean brightness is modulated, which, in principle, may influence the use of classical Cepheids for distance determination. Fortunately, the modulation of mean brightness does not exceed 0.01 mag in all but one star. Also, the effect averages out in typical observations spanning a long time base. Hence, in studies of sizeable samples of Cepheids with good phase coverage of the light curves, modulation has no practical effect on the distance determinations. It is clear however, that our understanding of the most important astrophysical candles is far from being complete.

is research is supported by the Polish National Science Center, ant DEC-2015/17/B/ST9/03421.

Importance of high-cadence and long-term monitoring



Fig. 3. Frequency spectrum for fundamental mode RR Lyr star (RRab) OGLE-BLG-RRLYR-01136. Spectra for OGLE-IV (top) and OGLE-BLG-RRLYR-01136. Spectra for OGLE-IV (top) the fundamental mode and its harmonics (dashed lines). Addi-tional signals are clearly detected in the OGLE-IV data (arrows), while in OGLE-III their detection is not possible due to much larger moise level, resulting from sparse data sampling. OGLE-IV data consist of 3092 data points gathered over 4 observing seasons, while OGLE-III data counts only 356 points (5 seasons). From Smolec et al. (2016).



Fig. 4. To detect long-period modulations, long-baseline of the data is nec-essary. Top panel illustrates the photometric data for first overtone RR Lyr star (RRe), OGL-FBLG-RRLYR-12659, gathered over 12 year during the OGL-FII and OGL-FV observations. Amplitude modulation is clear. Time-dependent Fourier analysis shows that both amplitude (middle panel) and pulsation phase (bottom panel) vary and are correlated. Modulation period is $\approx 1770d$. Adapted from Netzel, Smolec & OGLE (2017, in pres.)

RR Lyrae stars in NGC 6362



NGC 6362 (Fig. 7) is a nearby globular cluster that was extensively observed by the CASE project with the Swope telescope (Fig. 1) from 1995 to 2009. The top-quality photometry (see Fig. 8) enabled a detailed study of the dynamical properties of its 34 RR Lyr-type variables (Smolec et al., 2017). In many respects, this globular cluster and its RR Lyr sample are record holders: * The incidence rates of Blazhko phenomenon, 69% in RRab vari-

ables and 19% in RRc variables, are among the highest reported in the ground-based observations so far.

 \star 63% of RRc variables are in fact double-periodic and belong to the ${\rm RR}_{0.61}$ class. It is the highest incidence rate of this form of pulsation in the ground-based observations so far. Other important results are:

* Two stars previously classified as RRab, turned to be anomalous RRd variables. Their light curves, with clear signature of modulation, are plotted in Fig. 8.

* Very rare double-periodic modulations were detected in two stars



NATIONAL SCIENCE CENTRE